


ROMANIAN MATHEMATICAL MAGAZINE

In any ΔABC with p_a, p_b, p_c
 → Spieker cevians, the following relationship holds :

$$0 \leq \frac{p_a - w_a}{a} + \frac{p_b - w_b}{b} + \frac{p_c - w_c}{c} < \frac{4}{3}$$

Proposed by Mohamed Amine Ben Ajiba-Tanger-Morocco

Solution by Soumava Chakraborty-Kolkata-India



Let AS produced meet BC at X and $m(\angle BAX) = \alpha$ and $m(\angle CAX) = \beta$ (say)
 and inradius of $\Delta DEF = r'$ (say)

$$\text{Now, } 16[DEF]^2 = 2 \sum \left(\frac{a^2}{4} \right) \left(\frac{b^2}{4} \right) - \sum \frac{a^4}{16} = \frac{1}{16} \left(2 \sum a^2 b^2 - \sum a^4 \right) = \frac{16r^2 s^2}{16}$$

$$\Rightarrow [DEF] = \frac{rs}{4} \Rightarrow r' \left(\frac{\frac{a}{2} + \frac{b}{2} + \frac{c}{2}}{2} \right) = \frac{rs}{4} \Rightarrow r' = \frac{r}{2} \rightarrow (1)$$

$$\because \text{Spieker center is incenter of } \Delta DEF, \therefore m(\angle AFS) = B + \frac{C}{2} = \frac{2B + C}{2} = \frac{B + \pi - A}{2}$$

$$= \frac{\pi}{2} - \frac{A - B}{2} \text{ and } m(\angle AES) = C + \frac{B}{2} = \frac{\pi}{2} - \frac{A - C}{2} \rightarrow (2)$$

Via (1), (2) and using cosine law on ΔAFS and ΔAES , we arrive at :

$$AS^2 = \frac{r^2}{4 \sin^2 \frac{C}{2}} + \frac{c^2}{4} - \left(\frac{2r}{2 \sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A - B}{2}$$

$$= \frac{r^2}{4 \sin^2 \frac{B}{2}} + \frac{b^2}{4} - \left(\frac{2r}{2 \sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A - C}{2}$$

$$\Rightarrow 2AS^2 \stackrel{(i)}{=} \frac{r^2}{4 \sin^2 \frac{C}{2}} + \frac{c^2}{4} - \left(\frac{2r}{2 \sin \frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin \frac{A - B}{2} + \frac{r^2}{4 \sin^2 \frac{B}{2}} + \frac{b^2}{4}$$

$$- \left(\frac{2r}{2 \sin \frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin \frac{A - C}{2}$$

$$\begin{aligned}
 & \text{Now, } \left(\frac{2r}{2\sin\frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin\frac{A-B}{2} + \left(\frac{2r}{2\sin\frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin\frac{A-C}{2} \\
 &= \frac{r}{2} \left(4R\cos\frac{C}{2} \sin\frac{A-B}{2} + 4R\cos\frac{B}{2} \sin\frac{A-C}{2} \right) \\
 &= Rr \left(2\sin\frac{A+B}{2} \sin\frac{A-B}{2} + 2\sin\frac{A+C}{2} \sin\frac{A-C}{2} \right) \\
 &= Rr \left(1 - 2\sin^2\frac{B}{2} + 1 - 2\sin^2\frac{C}{2} - 2 \left(1 - 2\sin^2\frac{A}{2} \right) \right) \\
 &= 2Rr \left(\frac{2a(s-b)(s-c) - b(s-c)(s-a) - c(s-a)(s-b)}{abc} \right) \\
 &= \frac{Rr}{8Rrs} (2a^3 + (b+c)a^2 - 2a(b^2+c^2) - (b+c)(b-c)^2) \\
 &= \frac{4(b+c)bcs\sin^2\frac{A}{2} - 2a \cdot 2bcc\cos A}{8s} = \frac{bc \left((2s-a)\sin^2\frac{A}{2} - a \left(1 - 2\sin^2\frac{A}{2} \right) \right)}{2s} \\
 &= \frac{bc \left((2s+a)\sin^2\frac{A}{2} - a \right)}{2s} = \frac{(2s+a)(s-b)(s-c)}{2s} - 2Rr \\
 &\Rightarrow - \left(\frac{2r}{2\sin\frac{C}{2}} \right) \left(\frac{c}{2} \right) \sin\frac{A-B}{2} - \left(\frac{2r}{2\sin\frac{B}{2}} \right) \left(\frac{b}{2} \right) \sin\frac{A-C}{2} \\
 &\quad \stackrel{(*)}{=} \frac{-(2s+a)(s-b)(s-c)}{2s} + 2Rr \\
 & \text{Again, } \frac{r^2}{4\sin^2\frac{B}{2}} + \frac{r^2}{4\sin^2\frac{C}{2}} = \frac{r^2}{4} \left(\frac{ca}{(s-c)(s-a)} + \frac{ab}{(s-a)(s-b)} \right) \\
 &= \frac{r^2}{4r^2s} (ca(s-b) + ab(s-c)) = \frac{ab+ca}{4} - 2Rr \stackrel{(**)}{=} \frac{r^2}{4\sin^2\frac{B}{2}} + \frac{r^2}{4\sin^2\frac{C}{2}} \\
 & \text{(i), (*), (**)} \Rightarrow 2AS^2 = \frac{b^2+c^2+ab+ca}{4} - \frac{(2s+a)(s-b)(s-c)}{2s} \\
 &= \frac{(a+b+c)(b^2+c^2+ab+ca) - (2a+b+c)(c+a-b)(a+b-c)}{8s} \\
 &= \frac{b^3+c^3-abc+a(2b^2+2c^2-a^2)}{4s} \Rightarrow 2AS^2 \stackrel{(ii)}{=} \frac{b^3+c^3-abc+a(4m_a^2)}{4s} \\
 & \text{Via sine law on } \triangle AFS, \frac{r}{2\sin\frac{C}{2}\sin\alpha} = \frac{AS}{\cos\frac{A-B}{2}} = \frac{cAS}{(a+b)\sin\frac{C}{2}} \\
 &\Rightarrow c\sin\alpha \stackrel{(***)}{=} \frac{r(a+b)}{2AS} \text{ and via sine law on } \triangle AES, b\sin\beta \stackrel{****}{=} \frac{r(a+c)}{2AS} \\
 & \text{Now, } [BAX] + [BAX] = [ABC] \Rightarrow \frac{1}{2}p_a c\sin\alpha + \frac{1}{2}p_a b\sin\beta = rs \\
 & \quad \stackrel{\text{via (***) and (****)}}{\Rightarrow} \frac{p_a(a+b+a+c)}{4AS} = s \Rightarrow p_a = \frac{4s}{2s+a} AS \\
 &\Rightarrow p_a^2 \stackrel{\text{via (ii)}}{=} \frac{16s^2}{(2s+a)^2} \cdot \frac{b^3+c^3-abc+a(4m_a^2)}{8s}
 \end{aligned}$$

$$\therefore p_a^2 \stackrel{(\bullet)}{=} \frac{2s}{(2s+a)^2} (b^3 + c^3 - abc + a(4m_a^2))$$

$$\begin{aligned} \text{Now, } b^3 + c^3 - abc + a(4m_a^2) &= b^3 + c^3 - abc + a(2b^2 + 2c^2 - a^2) \\ &= (b+c)(b^2 - bc + c^2) + a(b^2 - bc + c^2) + a(b^2 + c^2 - a^2) \\ &= 2s(b^2 - bc + c^2) + a(b^2 - bc + c^2 + bc - a^2) \\ &= (2s+a)(b^2 - bc + c^2) + a\left(\frac{(b+c)^2 - (b-c)^2}{4} - a^2\right) \\ &= (2s+a)(b^2 - bc + c^2) + \frac{a(b+c+2a)(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a)(b^2 - bc + c^2) + \frac{a(2s-a+2a)(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a) \cdot \frac{4b^2 + 4c^2 - 4bc + a(b+c-2a)}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a). \end{aligned}$$

$$\frac{4(z+x)^2 + 4(x+y)^2 - 4(z+x)(x+y) + (y+z)((z+x) + (x+y) - 2(y+z))}{4}$$

$$\begin{aligned} & - \frac{a(b-c)^2}{4} \quad (a = y+z, b = z+x, c = x+y) \\ &= (2s+a) \cdot \frac{4x(x+y+z) + 2x(y+z) + 3(y-z)^2}{4} - \frac{a(b-c)^2}{4} \\ &= (2s+a) \left(s(s-a) + \frac{3}{4}(b-c)^2 + \frac{a(s-a)}{2} \right) - \frac{a(b-c)^2}{4} \\ &= (2s+a) \left(s(s-a) + \frac{3}{4}(b-c)^2 + \frac{a(s-a)}{2} \right) - \frac{(a+2s-2s)(b-c)^2}{4} \\ &= (2s+a) \left(s(s-a) + \frac{(b-c)^2}{2} + \frac{a(s-a)}{2} \right) + \frac{s(b-c)^2}{2} \end{aligned}$$

$$\therefore b^3 + c^3 - abc + a(4m_a^2) \stackrel{(\bullet\bullet)}{=} (2s+a) \left(\frac{(s-a)(2s+a)}{2} + \frac{(b-c)^2}{2} \right) + \frac{s(b-c)^2}{2}$$

$$\begin{aligned} \therefore (\bullet), (\bullet\bullet) \Rightarrow p_a^2 &= \frac{2s}{(2s+a)^2} \left(\frac{(s-a)(2s+a)^2}{2} + \frac{(2s+a)(b-c)^2}{2} + \frac{s(b-c)^2}{2} \right) \\ &= s(s-a) + (b-c)^2 \left(\left(\frac{s}{2s+a} \right)^2 + \frac{s}{2s+a} + \frac{1}{4} - \frac{1}{4} \right) \\ &= s(s-a) - \frac{(b-c)^2}{4} + (b-c)^2 \cdot \left(\frac{s}{2s+a} + \frac{1}{2} \right)^2 \\ &= s(s-a) + \frac{(b-c)^2}{4} \left(\frac{(4s+a)^2}{(2s+a)^2} - 1 \right) \\ &\Rightarrow p_a^2 \stackrel{(\bullet\bullet\bullet)}{=} s(s-a) + \frac{s(3s+a)(b-c)^2}{(2s+a)^2} \end{aligned}$$

$$\begin{aligned} \text{Again, } p_a - w_a &\leq \frac{2sa|b-c|}{4s^2 - a^2} \Leftrightarrow \frac{s(3s+a)(2s-a)^2 + s(s-a)(2s+a)^2}{(4s^2 - a^2)^2} \cdot (b-c)^2 \\ &\leq \frac{4s^2 a^2 (b-c)^2}{(4s^2 - a^2)^2} + \frac{4sa \cdot w_a \cdot |b-c|}{4s^2 - a^2} \end{aligned}$$

$$\begin{aligned}
 &\Leftrightarrow \frac{s(3s+a)(2s-a)^2 + s(s-a)(2s+a)^2 - 4s^2a^2}{(4s^2-a^2)^2} \cdot |b-c| \leq \frac{4sa \cdot w_a}{4s^2-a^2} \\
 &\quad (\because |b-c| \geq 0) \Leftrightarrow \frac{8s^2(2s+a)(s-a)}{4s^2-a^2} \cdot |b-c| \leq 4sa \cdot w_a \\
 &\Leftrightarrow \frac{4s^2(2s+a)^2(s-a)^2}{(4s^2-a^2)^2} \cdot (b-c)^2 \leq a^2 \left(s(s-a) - \frac{s(s-a)(b-c)^2}{(2s-a)^2} \right) \\
 &\quad \Leftrightarrow \frac{4s^2(s-a)^2 + a^2s(s-a)}{(2s-a)^2} \cdot (b-c)^2 \leq a^2s(s-a) \\
 &\quad \Leftrightarrow \frac{s(s-a)(4s^2-4sa+a^2)}{(2s-a)^2} \cdot (b-c)^2 \leq a^2s(s-a) \\
 &\Leftrightarrow s(s-a) \cdot (b-c)^2 \leq a^2s(s-a) \Leftrightarrow s(s-a)(a^2 - (b-c)^2) \stackrel{?}{\geq} 0 \rightarrow \text{true} \\
 &\quad \therefore p_a - w_a \leq \frac{2sa|b-c|}{4s^2-a^2} \text{ and analogs} \\
 &\Rightarrow \frac{p_a - w_a}{a} + \frac{p_b - w_b}{b} + \frac{p_c - w_c}{c} \leq \sum_{\text{cyc}} \frac{2s \cdot |b-c|}{4s^2-a^2} \stackrel{|b-c| < a \text{ and analogs}}{<} \\
 &\quad \frac{2s}{(\prod_{\text{cyc}}(2s-a)) \cdot (\prod_{\text{cyc}}(2s+a))} \cdot \sum_{\text{cyc}} (a(4s^2-b^2)(4s^2-c^2)) \\
 &= \frac{2s \left(2s \cdot 16s^4 - 4s^2(\sum_{\text{cyc}} ab^2 + \sum_{\text{cyc}} a^2b) + 4Rrs(s^2 + 4Rr + r^2) \right)}{(2s(s^2 + 2Rr + r^2)) \cdot (2s(9s^2 + 6Rr + r^2))} \\
 &= \frac{4s^2 \left(16s^4 - 4s^2(s^2 - 2Rr + r^2) + 2Rr(s^2 + 4Rr + r^2) \right)}{4s^2(s^2 + 2Rr + r^2)(9s^2 + 6Rr + r^2)} \\
 &= \frac{4s^2 \left(16s^4 - 4s^2(s^2 - 2Rr + r^2) + 2Rr(s^2 + 4Rr + r^2) \right)}{4s^2(s^2 + 2Rr + r^2)(9s^2 + 6Rr + r^2)} \stackrel{?}{<} \frac{4}{3} \\
 &\Leftrightarrow 2r \left((33R + 26r)s^2 + r(12R^2 + 13Rr + 2r^2) \right) \stackrel{?}{>} 0 \rightarrow \text{true} \\
 &\quad \therefore \frac{p_a - w_a}{a} + \frac{p_b - w_b}{b} + \frac{p_c - w_c}{c} < \frac{4}{3} \text{ and } p_a^2 - w_a^2 = \\
 &\quad \frac{s(3s+a)(2s-a)^2 + s(s-a)(2s+a)^2}{(4s^2-a^2)^2} \cdot (b-c)^2 \geq 0 \Rightarrow \frac{p_a - w_a}{a} \geq 0 \\
 &\quad \text{and analogs } \therefore \frac{p_a - w_a}{a} + \frac{p_b - w_b}{b} + \frac{p_c - w_c}{c} \geq 0 \text{ and so,} \\
 &\quad 0 \leq \frac{p_a - w_a}{a} + \frac{p_b - w_b}{b} + \frac{p_c - w_c}{c} < \frac{4}{3} \quad \forall \triangle ABC, \\
 &\quad \text{with equality iff } \triangle ABC \text{ is equilateral (QED)}
 \end{aligned}$$