

ROMANIAN MATHEMATICAL MAGAZINE

In $\triangle ABC$ the following relationship holds:

$$\sum_{cyc} \csc\left(\frac{A}{2}\right) \left(\sqrt{\left(\operatorname{ctg}^3\left(\frac{B}{2}\right) + \operatorname{ctg}^3\left(\frac{C}{2}\right)\right)} + \sqrt{\left(\operatorname{ctg}^3\left(\frac{A}{2}\right) + \operatorname{ctg}^3\left(\frac{C}{2}\right)\right)} \right) \geq 12\sqrt{6\sqrt{3}}$$

Proposed by Zaza Mzhavanadze-Georgia

Solution by Mirsadix Muzefferov-Azerbaijan

$$\text{Let : } LHS = LHS_1 + LHS_2 + LHS_3$$

$$LHS_1 = \csc\left(\frac{A}{2}\right) \left(\sqrt{\left(\operatorname{ctg}^3\left(\frac{B}{2}\right) + \operatorname{ctg}^3\left(\frac{C}{2}\right)\right)} + \sqrt{\left(\operatorname{ctg}^3\left(\frac{B}{2}\right) + \operatorname{ctg}^3\left(\frac{C}{2}\right)\right)} \right) \stackrel{AM-GM}{\geq}$$

$$\csc\left(\frac{A}{2}\right) \left(\sqrt{2\left(\operatorname{ctg}\left(\frac{B}{2}\right) \cdot \operatorname{ctg}\left(\frac{C}{2}\right)\right)^{\frac{3}{2}}} + \sqrt{2\left(\operatorname{ctg}\left(\frac{B}{2}\right) \cdot \operatorname{ctg}\left(\frac{C}{2}\right)\right)^{\frac{3}{2}}} \right) \stackrel{AM-GM}{\geq}$$

$$\csc\left(\frac{A}{2}\right) \left(\sqrt[4]{4\left(\operatorname{ctg}\left(\frac{A}{2}\right) \cdot \operatorname{ctg}\left(\frac{B}{2}\right) \cdot \operatorname{ctg}\left(\frac{C}{2}\right)\right)^{\frac{3}{2}} \cdot \left(\operatorname{ctg}\left(\frac{C}{2}\right)\right)^{\frac{3}{2}}} \right)$$

Analogously, it is calculated for LHS_2 and LHS_3

$$LHS \stackrel{A-G}{\geq} 3^3 \sqrt[3]{LHS_1 \cdot LHS_2 \cdot LHS_3} =$$

$$= 3 \cdot 2 \left(64 \left(\operatorname{ctg}\left(\frac{A}{2}\right) \cdot \operatorname{ctg}\left(\frac{B}{2}\right) \cdot \operatorname{ctg}\left(\frac{C}{2}\right) \right)^{\frac{9}{2}} \cdot \left(\operatorname{ctg}\left(\frac{A}{2}\right) \cdot \operatorname{ctg}\left(\frac{B}{2}\right) \cdot \operatorname{ctg}\left(\frac{C}{2}\right) \right)^{\frac{3}{2}} \right)^{\frac{1}{12}} \cdot$$

$$\left(\csc\left(\frac{A}{2}\right) \cdot \csc\left(\frac{B}{2}\right) \cdot \csc\left(\frac{C}{2}\right) \right)^{\frac{1}{3}} =$$

$$6 \left(2 \left(\operatorname{ctg}\left(\frac{A}{2}\right) \cdot \operatorname{ctg}\left(\frac{B}{2}\right) \cdot \operatorname{ctg}\left(\frac{C}{2}\right) \right) \right)^{\frac{1}{2}} \cdot \left(\csc\left(\frac{A}{2}\right) \cdot \csc\left(\frac{B}{2}\right) \cdot \csc\left(\frac{C}{2}\right) \right)^{\frac{1}{3}} =$$

$$6 \left(2 \left(\frac{p}{r} \right) \right)^{\frac{1}{2}} \cdot \left(\frac{4R}{r} \right)^{\frac{1}{3}} \stackrel{\text{Mitrinovic, Euler}}{\geq} 6(2 \cdot 3\sqrt{3})^{\frac{1}{2}} \cdot (8)^{\frac{1}{3}} = 12\sqrt{6\sqrt{3}}$$

Equality holds if : $A = B = C$