

ROMANIAN MATHEMATICAL MAGAZINE

In $\triangle ABC$ the following relationship holds:

$$\sum_{cyc} a \left(\tan \frac{A}{2} - \tan \frac{B}{2} \right) \geq 0$$

Proposed by Nguyen Hung Cuong – Vietnam

Solution by Daniel Sitaru – Romania

$$\text{WLOG: } a \leq b \leq c \Rightarrow \tan \frac{A}{2} \leq \tan \frac{B}{2} \leq \tan \frac{C}{2}$$

By rearrangements' inequality:

$$\begin{aligned} a \tan \frac{A}{2} + b \tan \frac{B}{2} + c \tan \frac{C}{2} &\geq a \tan \frac{B}{2} + b \tan \frac{C}{2} + c \tan \frac{A}{2} \\ a \left(\tan \frac{A}{2} - \tan \frac{B}{2} \right) + b \left(\tan \frac{B}{2} - \tan \frac{C}{2} \right) + c \left(\tan \frac{C}{2} - \tan \frac{A}{2} \right) &\geq 0 \end{aligned}$$

$$\sum_{cyc} a \left(\tan \frac{A}{2} - \tan \frac{B}{2} \right) \geq 0$$

Equality holds for $a = b = c$.